

CA125/MUC16 Catalog # PVGS1633

Product Information

Primary Accession Species	<u>Q8WXI7</u> Human
Sequence	Gly12660-Met12923
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	\leq 1 EU/ Ig of protein by LAL method
Biological Activity	Immobilized Human CA125, His & Avi Tag at 5.0 \Box /ml (100 \Box /Well). Dose response curve for Human MSLN with the EC ₅₀ of 3.78 \Box /ml determined by ELISA.
Expression System	Expi293
Formulation	Lyophilized from a 0.22 Im filtered solution in PBS, pH 7.4. Normally 5 % trehalose is added as protectant before lyophilization.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in distilled water up to 100 g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	94025
Other Names	Mucin-16, MUC-16, Ovarian cancer-related tumor marker CA125, CA-125, Ovarian carcinoma antigen CA125, MUC16 (<u>HGNC:15582</u>)
Target Background	MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy. Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces. Binding to MSLN mediates heterotypic cell adhesion. This may contribute to the metastasis of ovarian cancer to the peritoneum by initiating cell attachment to the mesothelial epithelium via binding to MSLN.

Protein Information

Name	MUC16 (<u>HGNC:15582</u>)
Function	Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces.
Cellular Location	Cell membrane; Single-pass type I membrane protein. Secreted, extracellular space. Note=May be liberated into the extracellular space following the phosphorylation of the intracellular C-terminus which induces the proteolytic cleavage and liberation of the extracellular domain
Tissue Location	Expressed in corneal and conjunctival epithelia (at protein level). Overexpressed in ovarian carcinomas and ovarian low malignant potential (LMP) tumors as compared to the expression in normal ovarian tissue and ovarian adenomas

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.