

# EpCAM/Trop1

Catalog # PVGS1639

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P16422</a> Human
<b>Sequence</b>	Gln24-Lys245
<b>Purity</b>	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
<b>Endotoxin Level</b>	≤ 1 EU/ $\mu$ g of protein by LAL method
<b>Biological Activity</b>	Immobilized Human EpCAM at 0.2 $\mu$ g/ml (100 $\mu$ l/Well). Dose response curve for Anti-Trop1 Ab with the EC <sub>50</sub> of 8.7 ng/ml determined by ELISA.
<b>Expression System</b>	Expi293
<b>Formulation</b>	Lyophilized from a 0.22 $\mu$ m filtered solution in PBS, pH 7.4. Normally 5 % trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in distilled water up to 100 $\mu$ g/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Gene ID</b>	4072
<b>Other Names</b>	Epithelial cell adhesion molecule, Ep-CAM, Adenocarcinoma-associated antigen, Cell surface glycoprotein Trop-1, Epithelial cell surface antigen, Epithelial glycoprotein, EGP, Epithelial glycoprotein 314, EGP314, hEGP314, KS 1/4 antigen, KSA, Major gastrointestinal tumor-associated protein GA733-2, Tumor-associated calcium signal transducer 1, CD326, EPCAM, GA733-2, M1S2, M4S1, MIC18, TACSTD1, TROP1
<b>Target Background</b>	Epithelial Cellular Adhesion Molecule (EpCAM), also known as KS1/4, gp40, GA733-2, 17-1A, and TROP-1, is a 40 kDa transmembrane glycoprotein that consists of a 242 amino acid (aa) extracellular domain with two EGF-like repeats, a 23 aa transmembrane segment, and a 26 aa cytoplasmic domain.

## Protein Information

---

<b>Name</b>	EPCAM
<b>Synonyms</b>	GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO
<b>Function</b>	May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.
<b>Cellular Location</b>	Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction. Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction
<b>Tissue Location</b>	Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.