

Spike Protein S1

Catalog # PVGS1643

Product Information

Primary Accession	P0DT2
Species	SARS-CoV-2
Sequence	Gln14-Arg685 (T95I, G142D, E154K, L452R, E484Q, D614G, P681R)
Purity	> 90% as analyzed by SDS-PAGE
Endotoxin Level	
Biological Activity	This protein is validated to bind with human ACE2 in functional ELISA assay.
Expression System	CHO
Theoretical Molecular Weight	75.7 kDa
Formulation	Supplied as a solution in PBS, pH 7.4.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -20°C or below. Please avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	43740568
Other Names	Spike glycoprotein {ECO:0000255 HAMAP-Rule:MF_04099}, S glycoprotein {ECO:0000255 HAMAP-Rule:MF_04099}, E2 {ECO:0000255 HAMAP-Rule:MF_04099}, Peplomer protein {ECO:0000255 HAMAP-Rule:MF_04099}, Spike protein S1 {ECO:0000255 HAMAP-Rule:MF_04099}, Spike protein S2 {ECO:0000255 HAMAP-Rule:MF_04099}, Spike protein S2' {ECO:0000255 HAMAP-Rule:MF_04099}, S {ECO:0000255 HAMAP-Rule:MF_04099}
Target Background	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as 2019-nCoV (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. As of May 2021, three sublineages have been found. Despite its name, B.1.617.3 was the first sublineage of this variant to be detected, in October 2020 in India. This sublineage has remained relatively uncommon compared to the two other sublineages, B.1.617.1 (also known as variant Kappa) and B.1.617.2 (also known as variant Delta), both of which were first detected in December 2020. This variant has the double mutations E484Q and L452R in the spike proteins. Emerging research suggests the variant may be more transmissible than previously evolved ones. Whether the effectiveness of currently-deployed vaccines is affected remains under investigation. Moreover, the sublineage B.1.617.2 has been redesignated as "variant of concern" (VOC-21APR-02) in May 2021, which

spreads more quickly than the original version of the virus.

Protein Information

Name	S {ECO:0000255 HAMAP-Rule:MF_04099}
Function	[Spike protein S1]: Attaches the virion to the cell membrane by interacting with host receptor, initiating the infection. The major receptor is host ACE2 (PubMed: 32142651 , PubMed: 32155444 , PubMed: 33607086). When S2/S2' has been cleaved, binding to the receptor triggers direct fusion at the cell membrane (PubMed: 34561887). When S2/S2' has not been cleaved, binding to the receptor results in internalization of the virus by endocytosis using host TFRC and GRM2 and leading to fusion of the virion membrane with the host endosomal membrane (PubMed: 32075877 , PubMed: 32221306 , PubMed: 34903715 , PubMed: 36779763). Alternatively, may use NRP1/NRP2 (PubMed: 33082294 , PubMed: 33082293) and integrin as entry receptors (PubMed: 35150743). The use of NRP1/NRP2 receptors may explain the tropism of the virus in human olfactory epithelial cells, which express these molecules at high levels but ACE2 at low levels (PubMed: 33082293). Uses also ASGR1 as an alternative receptor in an ACE2-independent manner (PubMed: 34837059). The stalk domain of S contains three hinges, giving the head unexpected orientational freedom (PubMed: 32817270).
Cellular Location	Virion membrane {ECO:0000255 HAMAP-Rule:MF_04099, ECO:0000269 PubMed:32979942}; Single-pass type I membrane protein {ECO:0000255 HAMAP-Rule:MF_04099, ECO:0000269 PubMed:34504087}, Host endoplasmic reticulum-Golgi intermediate compartment membrane {ECO:0000255 HAMAP-Rule:MF_04099, ECO:0000269 PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255 HAMAP-Rule:MF_04099}, Host cell membrane {ECO:0000255 HAMAP-Rule:MF_04099, ECO:0000269 PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255 HAMAP-Rule:MF_04099}. Note=Accumulates in the endoplasmic reticulum-Golgi intermediate compartment, where it participates in virus particle assembly. Some S oligomers are transported to the host plasma membrane, where they may mediate cell-cell fusion (PubMed:34504087). An average of 26 +/-15 S trimers are found randomly distributed at the surface of the virion (PubMed:32979942) {ECO:0000255 HAMAP-Rule:MF_04099, ECO:0000269 PubMed:32979942, ECO:0000269 PubMed:34504087}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.