

FGF-12

Catalog # PVGS1648

Product Information

Primary Accession P61328-2
Species Human

Sequence Met1-Thr181

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Biological Activity Immobilized Human FGF-12 at 2.0 ug/ml can bind Human FGFR3-Fc. The ED₅₀

of Human FGF-12 is 0.5-4.0 [g/ml.

Expression System E. coli

Formulation Lyophilized from a 0.2 Im filtered solution in 20 mM PB, 150 mM NaCl, 5 mM

EDTA, pH 7.5.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

distilled water up to 100 ☐g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at -70°C or

-20°C. Upon reconstitution, the product should be stable for up to 1 week at 4-7°C and up to 3 months at -20 °C or below. Avoid repeated freeze-thaw

cycles.

Additional Information

Target Background Fibroblast Growth Factor 12 (FGF-12) is a member of the fibroblast growth

factor (FGF) family. FGF-12 is probably involved in nervous system development and function. FGF-12 lacks the N-terminal signal sequence present in most of the FGF family members, but it contains clusters of basic residues that have been demonstrated to act as a nuclear localization signal. When transfected into mammalian cells, this protein accumulated in the nucleus, but was not secreted. The specific function of this gene has not yet been determined. Two alternatively spliced transcript variants encoding

distinct isoforms have been reported.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.