

CNTF

Catalog # PVGS1667

Product Information

Primary Accession P51642
Species Mouse

Sequence Ala2-Met198

Purity > 95% as analyzed by SDS-PAGE

> 95% as analyzed by HPLC

Endotoxin Level

Expression System E. coli

Theoretical Molecular Weight 22.5 kDa

Formulation Reconstitution

Lyophilized from a 0.2 $\,^{\circ}$ Im filtered solution in 2 × PBS, pH 7.4, 2 % trehalose. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a

concentration of 0.1 mg/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at -20°C or

-70°C. Upon reconstitution, the product should be stable for up to 1 week at

2-8°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID 12803

Other Names Ciliary neurotrophic factor, CNTF, Cntf

Target Background Ciliary neurotrophic factor (CNTF) is a polypeptide hormone whose actions

appear to be restricted to the nervous system where it promotes neurotransmitter synthesis and neurite outgrowth in certain neuronal populations. CNTF was initially identified as a trophic factor for embryonic chick ciliary parasympathetic neurons in culture. Furthermore, the protein is also a potent survival factor for neurons and oligodendrocytes and may be relevant in reducing tissue destruction during inflammatory attacks. In addition, CNTF is useful for treatment of motor neuron disease and it could reduce food intake without causing hunger or stress. Recombinant murine CNTF containing 198 amino acids and it shares 82 % and 95 % a.a. sequence

identity with human and rat CNTF.

Protein Information

Name Cntf

CNTF is a survival factor for various neuronal cell types. Seems to prevent the degeneration of motor axons after axotomy. **Function**

Cellular Location Cytoplasm.

Nervous system. **Tissue Location**

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.