

# Spike protein S1

Catalog # PVGS1681

## Product Information

|                                     |  |
|-------------------------------------|--|
| <b>Primary Accession Species</b>    | <a href="#">P0DTC2</a><br>SARS-CoV-2   |
| <b>Sequence</b>                     | Gln14-Arg685 (A67V, del69-70, T95I, G142D, del143-145, del211, L212I, ins214EPE, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H) |
| <b>Purity</b>                       | ≥ 90% as analyzed by SDS-PAGE  |
| <b>Endotoxin Level</b>              | ≤ 1 EU/ µg of protein by gel clotting method   |
| <b>Biological Activity</b>          | SARS-CoV-2 Spike protein S1, Omicron Variant, His Tag can bind with human ACE2 (Cat. No.: <a href="#">Z03516</a> ) in functional ELISA assay.  |
| <b>Expression System</b>            | 293 Cells  |
| <b>Theoretical Molecular Weight</b> | 79 kDa   |
| <b>Formulation</b>                  | Supplied as a solution in PBS pH 7.4   |
| <b>Storage &amp; Stability</b>      | Upon receiving, this product remains stable for up to 6 months at -20 °C or below. Avoid repeated freeze-thaw cycles.  |

## Additional Information

|                          |  |
|--------------------------|--|
| <b>Gene ID</b>           | 43740568   |
| <b>Other Names</b>       | Spike glycoprotein {ECO:0000255 HAMAP-Rule:MF_04099}, S glycoprotein {ECO:0000255 HAMAP-Rule:MF_04099}, E2 {ECO:0000255 HAMAP-Rule:MF_04099}, Peplomer protein {ECO:0000255 HAMAP-Rule:MF_04099}, Spike protein S1 {ECO:0000255 HAMAP-Rule:MF_04099}, Spike protein S2 {ECO:0000255 HAMAP-Rule:MF_04099}, Spike protein S2' {ECO:0000255 HAMAP-Rule:MF_04099}, S {ECO:0000255 HAMAP-Rule:MF_04099}   |
| <b>Target Background</b> | SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as 2019-nCoV (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. Recently, the new B.1.1.529 variant was confirmed in South Africa and preliminary evidence suggests an increased risk of reinfection with this variant. The B.1.1.529 variant was first reported to WHO on 24 November 2021 and WHO has designated this variant as a VOC (Variant of Concern), named Omicron. There are more than 30 mutations in the spike protein. |

## Protein Information

---

|                          |   |
|--------------------------|---|
| <b>Name</b>              | S {ECO:0000255   HAMAP-Rule:MF_04099}   |
| <b>Function</b>          | [Spike protein S1]: Attaches the virion to the cell membrane by interacting with host receptor, initiating the infection. The major receptor is host ACE2 (PubMed: <a href="#">32142651</a> , PubMed: <a href="#">32155444</a> , PubMed: <a href="#">33607086</a> ). When S2/S2' has been cleaved, binding to the receptor triggers direct fusion at the cell membrane (PubMed: <a href="#">34561887</a> ). When S2/S2' has not been cleaved, binding to the receptor results in internalization of the virus by endocytosis using host TFRC and GRM2 and leading to fusion of the virion membrane with the host endosomal membrane (PubMed: <a href="#">32075877</a> , PubMed: <a href="#">32221306</a> , PubMed: <a href="#">34903715</a> , PubMed: <a href="#">36779763</a> ). Alternatively, may use NRP1/NRP2 (PubMed: <a href="#">33082294</a> , PubMed: <a href="#">33082293</a> ) and integrin as entry receptors (PubMed: <a href="#">35150743</a> ). The use of NRP1/NRP2 receptors may explain the tropism of the virus in human olfactory epithelial cells, which express these molecules at high levels but ACE2 at low levels (PubMed: <a href="#">33082293</a> ). The stalk domain of S contains three hinges, giving the head unexpected orientational freedom (PubMed: <a href="#">32817270</a> ). |
| <b>Cellular Location</b> | Virion membrane {ECO:0000255   HAMAP-Rule:MF_04099, ECO:0000269   PubMed:32979942}; Single-pass type I membrane protein {ECO:0000255   HAMAP-Rule:MF_04099, ECO:0000269   PubMed:34504087}. Host endoplasmic reticulum-Golgi intermediate compartment membrane {ECO:0000255   HAMAP-Rule:MF_04099, ECO:0000269   PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255   HAMAP-Rule:MF_04099}. Host cell membrane {ECO:0000255   HAMAP-Rule:MF_04099, ECO:0000269   PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255   HAMAP-Rule:MF_04099}. Note=Accumulates in the endoplasmic reticulum-Golgi intermediate compartment, where it participates in virus particle assembly. Some S oligomers are transported to the host plasma membrane, where they may mediate cell-cell fusion (PubMed:34504087). An average of 26 +/-15 S trimers are found randomly distributed at the surface of the virion (PubMed:32979942) {ECO:0000255   HAMAP-Rule:MF_04099, ECO:0000269   PubMed:32979942, ECO:0000269   PubMed:34504087}   |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.