10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CEACAM-6/CD66c

Catalog # PVGS1715

Product Information

Primary Accession	P40199
Species	Human
Sequence	Lys35-Gly320
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per □g by the LAL method.
Expression System	HEK293
Theoretical Molecular Weight	34.1 kDa
Formulation	Lyophilized from a 0.22 □m filtered solution in PBS, pH 7.4.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 □g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	4680
Other Names	Cell adhesion molecule CEACAM6, Carcinoembryonic antigen-related cell adhesion molecule 6, CEA cell adhesion molecule 6 {ECO:0000312|HGNC:HGNC:1818}, Non-specific crossreacting antigen, Normal cross-reacting antigen, CD66c, CEACAM6 (HGNC:1818)
Target Background	CEACAM6 (carcinoembryonic antigen-related cell adhesion molecule 6), is encoded by the human CEACAM6 gene. It plays a probable role in both cell-cell signaling and cell adhesion, promoting tumor progression, invasion and metastasis. It is upregulated in pancreatic adenocarcinoma, breast cancer and non-small cell lung cancer.

Protein Information

Name	CEACAM6 (HGNC:1818)

Function	Cell surface glycoprotein that plays a role in cell adhesion and tumor progression (PubMed:10910050, PubMed:11590190, PubMed:1378450, PubMed:16204051, PubMed:2022629, PubMed:2803308, PubMed:8776764). Intercellular adhesion occurs in a calcium- and fibronectin-independent manner (PubMed:16204051, PubMed:2022629). Mediates homophilic and heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM5 and CEACAM8 (PubMed:11590190, PubMed:16204051, PubMed:2022629, PubMed:2803308, PubMed:8776764). Heterophilic interaction with CEACAM8 occurs in activated neutrophils (PubMed:8776764). Plays a role in neutrophil adhesion to cytokine-activated endothelial cells (PubMed:1378450). Plays a role in cell migration and cell adhesion to endothelial cells (PubMed:16204051).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor. Apical cell membrane. Cell surface. Note=Localized to the apical glycocalyx surface.
Tissue Location	Expressed in neutrophils (PubMed:1378450). Expressed in columnar epithelial and goblet cells of the colon (PubMed:10436421). Expressed in numerous tumor cell lines (at protein level) (PubMed:16204051).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.