

TROP-2/TACSTD2

Catalog # PVGS1717

Product Information

Primary Accession Species	P09758 Human
Sequence	His27-Thr274
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per μ g by the LAL method.
Biological Activity	Immobilized TROP-2/TACSTD2, His, Human (Cat.No.: Z03839) at 0.5 μ g/ml (100 μ l/Well) on the plate can bind Anti-TROP-2 Antibody.
Expression System	HEK293
Theoretical Molecular Weight	28.9 kDa
Formulation Reconstitution	Lyophilized from a 0.22 μ m filtered solution in PBS, pH 7.4. Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	4070
Other Names	Tumor-associated calcium signal transducer 2, Cell surface glycoprotein Trop-2, Membrane component chromosome 1 surface marker 1, Pancreatic carcinoma marker protein GA733-1, TACSTD2, GA733-1, M1S1, TROP2
Target Background	Trop-2 (Tumor-associated calcium signal transducer 2) is also known as epithelial glycoprotein-1 antigen (EGP-1). It is encoded by the TACSTD2 gene. The mutations of Trop-2 gene cause an autosomal recessive disorder. Trop-2 causes cancer cell growth, proliferation, invasion, migration, and survival of cancer cells

Protein Information

Name	TACSTD2
Synonyms	GA733-1, M1S1, TROP2
Function	May function as a growth factor receptor.
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Placenta, pancreatic carcinoma cell lines.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.