10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TROP-2/TACSTD2

Catalog # PVGS1721

Product Information

Primary Accession	Q8BGV3.1
Species	Mouse
Sequence	Gln25-Gly270
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per □g by the LAL method.
Expression System	HEK293
Theoretical Molecular Weight	28.8 kDa
Formulation	Lyophilized from a 0.22 □m filtered solution in PBS, pH 7.4.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 □g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	Trop-2 (Tumor-associated calcium signal transducer 2) is also known as epithelial glycoprotein-1 antigen (EGP-1). It is encoded by the TACSTD2 gene. The mutations of Trop-2 gene cause an autosomal recessive disorder. Trop-2 causes cancer cell growth, proliferation, invasion, migration, and survival of cancer cells

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.