

BCMA/TNFRSF17

Catalog # PVGS1725

Product Information

Primary Accession Species	Q02223-1 Human
Sequence	Met1-Ala54
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per μ g by the LAL method.
Biological Activity	Immobilized Human APRIL Trimer, His Tagat 0.1 μ g/ml (100 μ l/well) on the plate can bind BCMA/TNFRSF17 hFc Chimera, Human (Cat.No.: Z03847).
Expression System	HEK293
Theoretical Molecular Weight	31.4 kDa
Formulation Reconstitution	Lyophilized from a 0.22 μ m filtered solution in PBS, pH 7.4. Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	BCMA (B-cell maturation antigen), also called TNFRSF17, is encoded by the TNFRSF17 gene. It acts as a cell surface receptor of the TNF receptor superfamily. BCMA recognizes B-cell activating factor (BAFF). It interacts with the B-cell activating factor TNFSF13B.
--------------------------	--

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.