

# BCMA/TNFRSF17

Catalog # PVGS1728

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">A0A2K5UD97 / F7FFA0</a> Cynomolgus/Rhesus macaque
<b>Sequence</b>	Met1-Ala53
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
<b>Endotoxin Level</b>	Less than 1EU per $\mu$ g by the LAL method.
<b>Biological Activity</b>	Immobilized BCMA/TNFRSF17 , His, Cynomolgus/Rhesus macaque (Cat.No.: Z03852) at 1 $\mu$ g/ml (100 $\mu$ l/Well) on the plate can bind Biotinylated Human APRIL (Trimer), His Tag.
<b>Expression System</b>	HEK293
<b>Theoretical Molecular Weight</b>	7.10 kDa
<b>Formulation Reconstitution</b>	Lyophilized from a 0.22 $\mu$ m filtered solution in PBS, pH 7.4. Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage &amp; Stability</b>	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Target Background</b>	BCMA (B-cell maturation antigen), also called TNFRSF17, is encoded by the TNFRSF17 gene. It acts as a cell surface receptor of the TNF receptor superfamily. BCMA recognizes B-cell activating factor (BAFF). It interacts with the B-cell activating factor TNFSF13B.
--------------------------	--

## Protein Information

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.