

MSLN/Mesothelin

Catalog # PVGS1740

Product Information

Primary Accession Species	Q13421-3 Human
Sequence	Met487-Ser598
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per μ g by the LAL method.
Biological Activity	Immobilized MSLN/Mesothelin(487-598) hFc Chimera, Human (Cat.No.: Z03864) at 1 μ g/ml (100 μ l/Well) on the plate can bind Biotinylated Anti-MSLN Antibody.
Expression System	HEK293
Theoretical Molecular Weight	38.35 kDa
Formulation Reconstitution	Lyophilized from a 0.22 μ m filtered solution in PBS, pH 7.4. Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	Mesothelin, also known as MSLN, is encoded by the MSLN gene. It encodes a precursor protein of 71 kDa that is processed to a 31 kDa shed protein called megakaryocyte potentiating factor (MPF) and a 40 kDa fragment, mesothelin. MSLN is attached to the cell membrane by a glycosyl-phosphatidylinositol (GPI) anchor. It is a differentiation antigen which is highly expressed in several human cancers, including virtually all mesotheliomas and pancreatic adenocarcinomas.
--------------------------	---

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.