

MSLN/Mesothelin

Catalog # PVGS1750

Product Information

| | |
|---------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary Accession Species | Q61468-1 Mouse |
| Sequence | Asp298-Ser600 |
| Purity | > 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC |
| Endotoxin Level | Less than 1EU per μ g by the LAL method. |
| Biological Activity | Immobilized Human CA125, His Tag at 5 μ g/ml (100 μ l/Well) on the plate can bind MSLN/Mesothelin (298-600)[Biotin], His, Mouse (Cat.No.: Z03877). |
| Expression System | HEK293 |
| Theoretical Molecular Weight | 35.2 kDa |
| Formulation Reconstitution | Lyophilized from a 0.22 μ m filtered solution in PBS, pH 7.4. Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage & Stability | Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles. |

Additional Information

| | |
|--------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target Background | Mesothelin, also known as MSLN, is encoded by the MSLN gene. It encodes a precursor protein of 71 kDa that is processed to a 31 kDa shed protein called megakaryocyte potentiating factor (MPF) and a 40 kDa fragment, mesothelin. MSLN is attached to the cell membrane by a glycosyl-phosphatidylinositol (GPI) anchor. It is a differentiation antigen which is highly expressed in several human cancers, including virtually all mesotheliomas and pancreatic adenocarcinomas. |
|--------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.