

## B7-H3/CD276

Catalog # PVGS1785

## **Product Information**

Primary Accession Q8VE98
Species Mouse

Sequence Val29-Ala248.

**Purity** > 95% as determined by Bis-Tris PAGE

> 95% as determined by HPLC

**Endotoxin Level** Less than 1EU per g by the LAL method.

**Expression System** HEK293

Theoretical Molecular Weight 24.98 kDa

**Formulation** Lyophilized from a 0.22 Im filtered solution in PBS, pH 7.4.

**Reconstitution** It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH<sub>2</sub>O more than 100 □g/ml.

**Storage & Stability** Upon receiving, the product remains stable up to 6 months at -20 °C or below.

Upon reconstitution, the product should be stable for 3 months at -80 °C.

Avoid repeated freeze-thaw cycles.

## **Additional Information**

**Gene ID** 102657

Other Names CD276 antigen, B7 homolog 3, B7-H3, Costimulatory molecule, CD276, Cd276,

B7h3

**Target Background** B7-H3, a member of the B7 family of immunomodulatory molecules, is

overexpressed in a wide range of solid cancers.B7-H3 binds to activated T cells via an as yet unidentified receptor. In assays using sub-optimal amount so anti-CD3 stimulation, 2Ig  $\square$ H3 enhances T cell proliferation, T cell interferon-gamma (IFN-gamma) production, and cytotoxic T cells induction.

## **Protein Information**

Name Cd276

Synonyms B7h3

**Function** Modulates T-cell-mediated immune responses and the development of

acute and chronic transplant rejection. Plays a positive regulatory role in bone

formation and has a dual role in the bone- immune interface. Induces antitumor immunity as it activates both acquired and innate immunity leading to natural killer cell and CD8 T- cell dependent killing of tumor cells.

**Cellular Location** Membrane; Single-pass type I membrane protein

Tissue Location Ubiquitous...

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.