

EGFR/HER1

Catalog # PVGS1789

Product Information

Primary Accession Species	P55245 Rhesus macaque
Sequence	Leu25-Ser645
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per μ g by the LAL method.
Biological Activity	Immobilized EGFR/HER1, His, Rhesus macaque (Cat.No.: Z03923) at 5 μ g/ml (100 μ l/Well) on the plate can bind Human EGF, hFc Tag
Expression System	HEK293
Theoretical Molecular Weight	69.8 kDa
Formulation	Lyophilized from a 0.22 μ m filtered solution in PBS, pH 7.4 .
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O more than 100 μ g/ml.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	613027
Other Names	Epidermal growth factor receptor, 2.7.10.1, EGFR
Target Background	The epidermal growth factor receptor is a transmembrane protein that is a receptor for members of the epidermal growth factor family of extracellular protein ligands. The epidermal growth factor receptor is a member of the ErbB family of receptors, a subfamily of four closely related receptor tyrosine kinases: EGFR, HER2/neu, Her 3 and Her 4. Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses.

Protein Information

Name	EGFR
Function	Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses. Known ligands include EGF, TGFA/TGF-alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling. Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (By similarity). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (By similarity). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P00533}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P00533} Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:P00533}; Single- pass type I membrane protein {ECO:0000250 UniProtKB:P00533}. Golgi apparatus membrane {ECO:0000250 UniProtKB:P00533}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P00533}. Nucleus membrane {ECO:0000250 UniProtKB:P00533}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P00533}. Endosome {ECO:0000250 UniProtKB:P00533}. Endosome membrane {ECO:0000250 UniProtKB:P00533}. Nucleus {ECO:0000250 UniProtKB:P00533} Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER. Endocytosed upon activation by ligand Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF). {ECO:0000250 UniProtKB:P00533}
Tissue Location	Hypothalamus..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.