

CD7

Catalog # PVGS1889

Product Information

Primary Accession Species	P50283 Mouse
Sequence	Gln24-Pro150
Purity	> 95% as determined by Bis-Tris PAGE
Endotoxin Level	Less than 1EU per μ g by the LAL method.
Biological Activity	Human SECTM1, hFc Tag captured on CM5 Chip via Protein A can bind CD7, His, Mouse in SPR assay (Biacore T200). Test result was comparable to standard batch.
Expression System	HEK293
Theoretical Molecular Weight	15.3 kDa
Formulation Reconstitution	Lyophilized from a 0.22 μ m filtered solution in PBS , (pH 7.4). Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	12516
Other Names	T-cell antigen CD7, CD7, Cd7
Target Background	CD7, also known as Leu-9, is an approximately 40 kDa glycosylated and palmitoylated transmembrane protein in the immunoglobulin superfamily. CD7 is expressed on T cells, NK cells , myeloid progenitor cells, and CD19 B progenitor cells. Among CD8 T cells, the CD7-bright population preferentially contains na $\bar{\nu}$ e and memory cells, while more weak expressors are primarily effector cells.

Protein Information

Name	Cd7
-------------	-----

Function	Transmembrane glycoprotein expressed by T-cells and natural killer (NK) cells and their precursors. Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1. In turn, mediates the production of cytokines such as IL-2. On resting NK-cells, CD7 activation results in a significant induction of gamma-interferon levels.
Cellular Location	Membrane; Single-pass type I membrane protein.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.