

CD3E&CD3D

Catalog # PVGS1897

Product Information

Primary Accession Species	P07766(CD3E)&P04234(CD3D) Human
Sequence	Asp23-Asp126(CD3E)&Phe22-Ala105(CD3D)
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per μ g by the LAL method.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized CD3E&CD3D hFc Chimera, Human at 2 μ g/ml (100 μ l/well) on the plate can bind Anti-CD3E&CD3D Ab.2, mFc Tag. Test result was comparable to standard batch.
Expression System	HEK293
Theoretical Molecular Weight	37.8 kDa (CD3E) and 35.4 kDa (CD3D)
Formulation Reconstitution	Lyophilized from a 0.22 μ m filtered solution in PBS , (pH 7.4). Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	T-cell surface glycoprotein CD3 epsilon & CD3 delta chain, also known as CD3E & CD3D , are single-pass type I membrane proteins. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain.
--------------------------	--

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.