

## Fc gamma RIV/CD16-2

Catalog # PVGS1901

## **Product Information**

Primary Accession <u>A0A0B4J1G0-1</u>

**Species** Mouse

**Sequence** Gly21-Gln203

**Purity** > 95% as determined by Bis-Tris PAGE

> 95% as determined by HPLC

**Endotoxin Level** Less than 1EU per Ig by the LAL method.

**Biological Activity** Fc gamma RIV/CD16-2, His, Mouse captured on CM5 Chip via antihis antibody

can bind AntiCD3 Antibody in SPR assay (Biacore T200). Test result was

comparable to standard batch.

Expression System HEK293

Theoretical Molecular Weight 22 kDa

**Formulation** Lyophilized from a 0.22 Im filtered solution in PBS, (pH 7.4).

**Reconstitution** Centrifuge the tube before opening. Reconstituting to a concentration more

than 100 ½/ml is recommended. Dissolve the lyophilized protein in distilled

water.

**Storage & Stability** Upon receiving, the product remains stable up to 6 months at -20 °C or below.

Upon reconstitution, the product should be stable for 3 months at -80 °C.

Avoid repeated freeze-thaw cycles.

## **Additional Information**

**Target Background** FcgammaRIV is a relatively new IgG Fc receptor (FcgammaR) that is reported

to contribute to the pathogenesis of autoimmune diseases. FcgammaRIII and FcgammaRIV are each essential to trigger an FcRgamma-linker for activation of T-cell-dependent signal that drives C5a production in the Arthus reaction. A combined requirement for FcgammaRIII and FcgammaRIV in autoimmune injury, and identify the linker for activation of T cells adaptor as an integral component of linked FcgammaR and C5a anaphylatoxin receptor activation to

generate inflammation.

## **Protein Information**

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.