

# Fc gamma RIII/CD16

Catalog # PVGS1902

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">Q5D5I8</a> Mouse
<b>Sequence</b>	Leu32-Thr215
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
<b>Endotoxin Level</b>	Less than 1EU per $\mu$ g by the LAL method.
<b>Biological Activity</b>	OKT3 captured on CM5 Chip via Protein A can bind Fc gamma RIII/CD16, His, Mouse in SPR assay (Biacore T200). Test result was comparable to standard batch.
<b>Expression System</b>	HEK293
<b>Theoretical Molecular Weight</b>	22.2 kDa
<b>Formulation Reconstitution</b>	Lyophilized from a 0.22 $\mu$ m filtered solution in PBS, (pH 7.4). Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage &amp; Stability</b>	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Target Background</b>	Immunoglobulin G (IgG) Fc receptors play a critical role in linking IgG antibody-mediated immune responses with cellular effector functions. A high resolution map of the binding site on human IgG1 for human Fc gamma RI, Fc gamma RIIA, Fc gamma RIIB, Fc gamma RIIIA, and FcRn receptors has been determined. A common set of IgG1 residues is involved in binding to all Fc gamma R; Fc gamma RII and Fc gamma RIII also utilize residues outside this common set.
--------------------------	---

## Protein Information

---