

# CD21

Catalog # PVGS1937

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P20023-1</a> Human
<b>Sequence</b>	Ile21-Arg971
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
<b>Endotoxin Level</b>	Less than 1EU per $\mu$ g by the LAL method.
<b>Biological Activity</b>	Measured by its binding ability in a functional ELISA. Immobilized CD21, His, Human at 1 $\mu$ g/ml (100 $\mu$ l/well) on the plate can bind Anti-CD21 Antibody, hFc Tag. Test result was comparable to standard batch.
<b>Expression System</b>	HEK293
<b>Theoretical Molecular Weight</b>	105.3 kDa
<b>Formulation Reconstitution</b>	Lyophilized from a 0.22 $\mu$ m filtered solution in PBS, (pH 7.4). Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage &amp; Stability</b>	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Target Background</b>	CD21 mRNA expression, an EBV-entry receptor, was transiently detected in NK cells after exosome treatment, and levels decreased after further culture. CD21 protein expression was also transiently transferred to NK cells after co-culture with an EBV-positive Burkitt lymphoma cell line (Raji) via trogocytosis. However, EBV did not infect NK cells through CD21-mediated trogocytosis.
--------------------------	--

## Protein Information

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.