10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Fc gamma RIIIA/CD16a (F176)

Catalog # PVGS1944

Product Information

Primary Accession	P08637-1
Species	Human
Sequence	Gly17-Gln208(F176)
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per μg by the LAL method.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized Fc gamma RIIIA/CD16a (F176), His & Avi, Human at 2 μg/ml (100 μl/well) on the plate can bind Anti-Fc gamma RIIIA Antibody, hFc Tag. Test result was comparable to standard batch.
Expression System	HEK293
Theoretical Molecular Weight	24.7 kDa
Formulation	Lyophilized from a 0.22 μm filtered solution in PBS, (pH 7.4).
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	Human Fc gamma RIIIA/CD16a Protein is a receptor for the Fc region of IgG. Binds complexed or aggregated IgG and also monomeric IgG. Mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.