

SARS-CoV-2 Nucleocapsid protein (His Tag)

Catalog # PVGS1950

Product Information

Primary Accession Species	P0DTC9 SARS-CoV-2
Sequence	Ser2-Ala419
Purity	> 90% as analyzed by SDS-PAGE
Biological Activity	SARS-CoV-2 Nucleocapsid protein (His Tag) can bind with SARS-CoV-2 Nucleocapsid Antibody (HC2003), Human Chimeric (Cat. No. A02039) in functional ELISA assay.
Expression System	E. coli
Theoretical Molecular Weight	46 kDa
Formulation Storage & Stability	Supplied as a solution in PBS pH 7.4 containing 10% glycerol. Upon receiving, this product remains stable for up to 6 months at -20°C or below. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	43740575
Other Names	Nucleoprotein {ECO:0000255 HAMAP-Rule:MF_04096}, N, Nucleocapsid protein {ECO:0000255 HAMAP-Rule:MF_04096}, NC {ECO:0000255 HAMAP-Rule:MF_04096}, Protein N {ECO:0000255 HAMAP-Rule:MF_04096}, N {ECO:0000255 HAMAP-Rule:MF_04096}
Target Background	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as 2019-nCoV (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. SARS-CoV-2 Nucleocapsid Protein is associated with nucleic acid. It is the most abundant protein for coronavirus. Because of the strong immunogenicity of coronavirus Nucleocapsid, it is believed that SARS-CoV-2 Nucleocapsid Protein has potential value for the diagnosis of the virus.

Protein Information

Name	N {ECO:0000255 HAMAP-Rule:MF_04096}
Function	Packages the positive strand viral genome RNA into a helical

ribonucleocapsid (RNP) and plays a fundamental role during virion assembly through its interactions with the viral genome and membrane protein M (PubMed:[33264373](#)). Plays an important role in enhancing the efficiency of subgenomic viral RNA transcription as well as viral replication. Attenuates the stress granules formation by reducing host G3BP1 access to host mRNAs under stress conditions (PubMed:[34901782](#), PubMed:[36534661](#)).

Cellular Location

Virion {ECO:0000255|HAMAP-Rule:MF_04096}. Host cytoplasm Secreted. Host extracellular space. Note=Probably associates with ER-derived membranes where it participates in viral RNA synthesis and virus budding. When located inside the virion, complexed with the viral RNA Can be secreted by unconventional protein secretion (UPS) (PubMed:35921414). When secreted, can bind to host glycosaminoglycans on infected and non infected cells (PubMed:35921414). Found in host cytoplasmic stress granules (PubMed:34901782). {ECO:0000255|HAMAP- Rule:MF_04096, ECO:0000269|PubMed:34901782, ECO:0000269|PubMed:35921414}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.