

# Heat Stable FGF-basic, Human

Catalog # PVGS1964

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P09038</a> Human
<b>Sequence</b>	Expressed with an N-terminal Met and several site mutations. Ala135-Ser288
<b>Purity</b>	≥ 95% as analyzed by SDS-PAGE
<b>Endotoxin Level Biological Activity</b>	ED <sub>50</sub> 50, the calculated specific activity is approximately $> 4.0 \times 10^6$ units/mg. It is recommended to experimentally determine the optimal concentration for each specific application by performing a dose response assay.
<b>Expression System</b>	E.coli
<b>Theoretical Molecular Weight</b>	17.1 kDa
<b>Formulation Reconstitution</b>	Lyophilized after extensive dialysis against PBS. Before opening, centrifuge the vial briefly to bring the contents to the bottom. Reconstitute the lyophilized powder in PBS up to 100 µg/ml
<b>Storage &amp; Stability</b>	Upon receiving, the lyophilized product remains stable for up to 6 months at lower than -70 °C. Upon reconstitution, the product is stable for up to 1 week at 4 °C or up to 3 months at -20 °C. Avoid repeated freeze-thaw cycles by making single-use aliquots before the solution is stored at -20 °C.

## Additional Information

---

<b>Gene ID</b>	2247
<b>Other Names</b>	Fibroblast growth factor 2, FGF-2, Basic fibroblast growth factor, bFGF, Heparin-binding growth factor 2, HBGF-2, FGF2, FGFB
<b>Target Background</b>	Heat Stable FGF-basic, Human is a pleiotropic cytokine and one of the prototypic members of the heparin-binding FGF family. Like other FGF family members, FGF-basic has the β trefoil structure. In vivo, FGF-basic is produced by a variety of cells, including cardiomyocytes, fibroblasts, and vascular cells. FGF-basic regulates a variety of processes including cell proliferation, differentiation, survival, adhesion, motility, apoptosis, limb formation and wound healing. FGF-basic can be tumorigenic due to its role in angiogenesis and blood vessel remodeling. The angiogenic effects of FGF-basic can produce beneficial cardioprotection during acute heart injury.

## Protein Information

---

<b>Name</b>	FGF2
<b>Synonyms</b>	FGFB
<b>Function</b>	Acts as a ligand for FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed: <a href="#">8663044</a> ). Also acts as an integrin ligand which is required for FGF2 signaling (PubMed: <a href="#">28302677</a> ). Binds to integrin ITGAV:ITGB3 (PubMed: <a href="#">28302677</a> ). Plays an important role in the regulation of cell survival, cell division, cell differentiation and cell migration (PubMed: <a href="#">28302677</a> , PubMed: <a href="#">8663044</a> ). Functions as a potent mitogen in vitro (PubMed: <a href="#">1721615</a> , PubMed: <a href="#">3732516</a> , PubMed: <a href="#">3964259</a> ). Can induce angiogenesis (PubMed: <a href="#">23469107</a> , PubMed: <a href="#">28302677</a> ). Mediates phosphorylation of ERK1/2 and thereby promotes retinal lens fiber differentiation (PubMed: <a href="#">29501879</a> ).
<b>Cellular Location</b>	Secreted. Nucleus. Note=Exported from cells by an endoplasmic reticulum (ER)/Golgi-independent mechanism. Unconventional secretion of FGF2 occurs by direct translocation across the plasma membrane (PubMed:20230531). Binding of exogenous FGF2 to FGFR facilitates endocytosis followed by translocation of FGF2 across endosomal membrane into the cytosol (PubMed:22321063). Nuclear import from the cytosol requires the classical nuclear import machinery, involving proteins KPNA1 and KPNB1, as well as CEP57 (PubMed:22321063)
<b>Tissue Location</b>	Expressed in granulosa and cumulus cells. Expressed in hepatocellular carcinoma cells, but not in non-cancerous liver tissue.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.